

The velocity and angular momentum of a free Dirac electron

Lu Lin

Department of Electrophysics
National Chiao Tung University
Hsinchu, Taiwan
Republic of China

ABSTRACT

It is shown that, in Dirac theory, there is a spatial velocity of a free electron which commutes with the Hamiltonian, so it is a conserved quantity of the motion. Furthermore, there is a spatial orbital angular momentum which also commutes with the Hamiltonian and is a constant of the motion.

1. The spatial velocity for a of Dirac electron:

To describe the motion of a free electron, besides the space-time coordinates $(\mathbf{X}, ct) = (x_1 x_2 x_3 x_4)$, Dirac introduced also $(\alpha_1 \alpha_2 \alpha_3 \beta) = (\boldsymbol{\alpha} \beta)$ for the internal motion in the electron which operate on a complex spinor space. Since the spatial space and the spinor space are basically independent, so the α 's and β are independent of and commute with the x 's and the p 's, where $(p_1 p_2 p_3 p_4)$ are the momentum and energy of the electron. The Hamiltonian of the free electron is given as

$$H = c\boldsymbol{\alpha} \cdot \mathbf{P} + mc^2\beta. \quad (1)$$

Dirac showed that

$$\frac{d}{dt}\mathbf{X} = c\boldsymbol{\alpha}, \quad (2)$$

$$\mathbf{X} = \boldsymbol{\xi} + \boldsymbol{\eta}, \quad (3)$$

$$\boldsymbol{\xi} = \boldsymbol{\xi}_0 + c^2\mathbf{P}H^{-1}t, \quad (4)$$

$$\boldsymbol{\eta} = \frac{i}{2}c\hbar(\boldsymbol{\alpha}H^{-1} - c\mathbf{P}H^{-2}) \quad (5)$$

where $\boldsymbol{\xi}_0$ is the $\boldsymbol{\xi}(t)$ when $t = 0$, and the σ 's the Pauli matrices. Next, we calculate the time derivative of $\boldsymbol{\eta}$ by taking the commutator of $\boldsymbol{\eta}$ with H and obtain

$$\frac{d}{dt}\boldsymbol{\eta} = c\boldsymbol{\alpha} - c^2\mathbf{P}H^{-1}. \quad (6)$$

From equation (2) and equation (6), we get

$$\frac{d}{dt}\boldsymbol{\xi} = \frac{d}{dt}(\mathbf{X} - \boldsymbol{\eta}) = c^2\mathbf{P}H^{-1}. \quad (7)$$

The operator $c^2 \mathbf{P} H^{-1}$ commutes with the Hamiltonian H , and has eigen value $\mathbf{v} = \mathbf{P}/M$ (M is the total mass), which is a velocity vector of a free particle. So it can be simultaneously measured with H and is a conserved quantity during the course motion. Therefore we conclude that $c^2 P_i H^{-1} = v_i$ is the eigen value obtained by a measurement of the i -component of the velocity of a free electron.

It may be worthwhile to mention that, since for the moment we do not know exactly the operation rules in the theory for the vector $\boldsymbol{\xi}$. So when we have to deal with this vector, we should express it as $\xi_i = x_i - \eta_i$, and then deal with x_i and η_i . Otherwise we may have trouble. For instance, the commutator for p and q is $[q, p] = i\hbar$, so $[x_1, p_1] = i\hbar = [\xi_1 + \eta_1, p_1] = [\xi_1, p_1] + [\eta_1, p_1]$. From equation (5) we see that η_1 commutes with p_1 so $[\eta_1, p_1] = 0$, and $[\xi, p_1] = i\hbar$. Then if we take $[x_1, H] = [\xi_1, H] + [\eta_1, H] = i\hbar \dot{x}_1$, we will get $\dot{x}_1 = c\alpha_1 + c\alpha_1 - c^2 p_1 H^{-1} = 2c\alpha_1 - c^2 p_1 H^{-1}$, which is wild when we compare it with equation (2). This example says that we can not operate directly with the $\xi's$. We have to operate with the $(x's - \eta's)$.

Consider two classical harmonic oscillators with coordinates z_1 and z_2 and their Hamiltonians H_1 and H_2 . When there is no coupling between them, z_1 and z_2 will oscillate independently and separately. However, if there is a coupling between them, the total Hamiltonian will become $H_1 + H_2 + H_{12}$. When we diagonalize the system, the coordinates z_1 and z_2 will get mixed, and the normal coordinates will be a mixture of z_1 and z_2 .

If Ψ is an eigenfunction of the Hamiltonian H in equation (1), then it is an eigenfunction of the Hamiltonian $H' = H^2$. The Hamiltonian H' has only spatial quantities (without α and β). Thus Ψ must be able to describe a system which has only spatial variables. On the other hand, H is a Hamiltonian which contains interactions between momentum and spin. So the generalized coordinates of the system (the x'_i s), which are the dynamical conjugate variables of momentum, must be a mixture of both spatial and spin variables as we can see from equations (3,4,5). The time derivatives of the x'_i s are then mixtures of the time derivatives of both kinds of variables. However, only the component which is pure spatial is experimentally measurable.

2. The spatial orbital angular momentum of a free Dirac electron:

Consider an angular momentum ℓ defined as

$$\ell = (\mathbf{X} - \boldsymbol{\eta}) \times \mathbf{P} = \boldsymbol{\xi} \times \mathbf{P}. \quad (8)$$

Since $\mathbf{P}$ is a constant vector of the motion, from equation (7) we see that the time derivative of ℓ vanishes. So ℓ commutes with H and is a constant of the motion and can be simultaneously measured with the Hamiltonian. Also, v_i is a spatial velocity and is a constant of motion. ξ_i is the time integration of v_i , so it is a spatial variable. Therefore, ℓ is the spatial angular momentum of the free electron.

It is well known that the electron has an intrinsic angular momentum of $\frac{\hbar}{2}\boldsymbol{\sigma}$ due to the $SU(2)$ symmetry. In reference (Dirac 1958), Dirac showed that the σ' s do not commute with the Hamiltonian (1), so the spin angular momentum is not a constant

of the motion. But the Hamiltonian (1) is rotational invariant, the total angular momentum must be conserved. Through the interaction between the spin and the spatial variables, the rotational symmetry will induce an angular momentum term, which arises from coupling the two parts of variables in order to conserve the total angular momentum. The simplest form of this term which meets those requirements is $\boldsymbol{\eta} \times \boldsymbol{P}$. Finally we have for the total angular momentum as

$$\boldsymbol{j} = \boldsymbol{\xi} \times \boldsymbol{P} + \boldsymbol{\eta} \times \boldsymbol{P} + \frac{\hbar}{2} \boldsymbol{\sigma}. \quad (9)$$

If we combine the first two terms together, we will just have $\boldsymbol{j} = \boldsymbol{X} \times \boldsymbol{P} + \frac{\hbar}{2} \boldsymbol{\sigma}$. However, equation (9) states that the total angular momentum is a sum of a spin, a spatial, and a spin-spatial coupling term.

The author wishes to thank Prof. Y.S. Wu for valuable discussions.

Reference

Dirac PAM 1958 Dirac PAM 1958 The principles of quantum mechanics, fourth ed. Oxford Univ. Press, London.